

Contents

INTRODUCTION

Rolf Hårdter and Thomas Fairhurst 1

BOTANICAL ASPECTS OF THE OIL PALM RELEVANT TO CROP MANAGEMENT

Ng Siew Kee, Helmut von Uexküll and Rolf Hårdter 13

LAND SELECTION FOR OIL PALM

S. Paramanathan 27

THE SEARCH FOR YIELD IN OIL PALM: BASIC PRINCIPLES

Kees Breure 59

CLONAL OIL PALM: PRODUCTION, YIELD PERFORMANCE AND NUTRITIONAL REQUIREMENTS

Ng Siew Kee, Thong Kar Cheong , Khaw Cheng Haw , Henry S.H. Ooi, Leng KaYee and P. Kayaroganam, Helmut von Uexküll and Rolf Hårdter 99

STANDARD AGRONOMIC PROCEDURES AND PRACTICES

Roger A. Gillbanks 115

LEGUME COVER PLANTS

Ken E. Giller and Thomas Fairhurst 151

CANOPY MANAGEMENT TO OPTIMIZE YIELD

Helmut von Uexküll, Ian E. Henson and Thomas Fairhurst 163

THE ROLE OF NUTRITION IN DISEASE CONTROL

Peter Turner 181

GENERAL OIL PALM NUTRITION

Kah-Joo Goh and Rolf Hårdter 191

ASSESSMENT OF OIL PALM FERTILIZER REQUIREMENTS

Hugh Foster 231

DESIGN AND ANALYSIS OF FERTILIZER EXPERIMENTS

Rob Verdooren 259

FERTILIZING FOR MAXIMUM RETURN

Kah-Joo Goh, Rolf Hårdter and Thomas Fairhurst 279

<i>UTILIZATION OF FIELD RESIDUES AND MILL BY-PRODUCTS</i>	
Mike Redshaw	307
<i>A CONCEPTUAL FRAMEWORK FOR PRECISION AGRICULTURE IN OIL PALM PLANTATIONS</i>	
Thomas Fairhurst, Ian Rankine, Armin Gfroerer Kerstan, Vince McAleer, Clive Taylor and William Griffiths	321
<i>CONCLUSIONS: FROM AREA EXPANSION TO YIELD INTENSIFICATION</i>	
Thomas Fairhurst and Rolf Härdter	333
<i>ALPHABETICAL INDEX</i>	369

Appendices

1	<i>GROUPING PLANTATION FIELDS IN LEAF SAMPLING UNITS</i>	338
2	<i>MARKING PALMS FOR LEAF SAMPLING IN LEAF SAMPLING UNITS</i>	339
3	<i>LEAF (FROND) SAMPLING</i>	340
4	<i>SOIL SAMPLING FOR ANALYSIS AND SOIL FERTILITY APPRAISAL</i>	342
5	<i>ISOMETRIC PALM POINT MAPPING</i>	344
6	<i>MEASUREMENT OF VEGETATIVE AND GENERATIVE GROWTH IN MATURE PALMS</i>	345
7	<i>MEASUREMENT OF NUTRIENT USE EFFICIENCY</i>	349
8	<i>SAMPLING FERTILIZER MATERIALS FOR ANALYSIS</i>	351

Color Plates

1	(a) <i>IKONOS satellite image showing delineation of field and natural reserve land at PT Harapan Sawit Lestari, Kalimantan, Indonesia. Areas bordering creeks have not been planted</i>	354
	(b) <i>Each planted field is referenced to a weather station. Climatic data is stored in the database management system which also calculates water deficits (data from PT Asiatic Persada, Jambi province, Indonesia)</i>	354
2	<i>Map showing yield gaps (attainable yield - actual yield) in 2002 in PT Asiatic Persada, an oil palm plantation in Jambi province, Indonesia, currently under rehabilitation</i> ..	355

3	Leaf phosphorus (P) levels portrayed in a discrete variability map showing deficient fields (red) and fields where leaf P status is above the critical threshold (green) in 2002 in PT Asiatic Persada, an oil palm plantation in Jambi province, Indonesia, currently under rehabilitation	356
4	(a) The database management system (DBMS) is used to compare leaf nutrient levels in different planting materials. Data can also be compared by year, unit, LSU group, and tree age (data of PT Harapan Sawit Lestari, West Kalimantan, Indonesia)	357
	(b) Qualitative environmental data and quantitative data on pest and disease incidence is stored and analyzed in the database management system (DBMS) (data of Milne Bay Estates, Milne Bay province, Papua New Guinea).....	357
5	Maps showing field erosion status by plotting the mode value for each field (a) or by using geographical information software to interpolate data collected at each LSU point to produce a continuous variability map (b) (data from PT Asiatic Persada, Jambi province, Indonesia).....	358
6	Isometric maps showing palm points in relation to block boundaries and waterways can be produced using GIS software and palm point data collected with a GPS	359
7	(a) A nitrogen (N) deficient palm growing on a well-drained soil	360
	(b) A palm affected by poor drainage. The ability of soil to supply nitrogen is reduced under very wet conditions.....	360
	(c) Vegetative growth is reduced and fronds are shorter in palms affected by phosphorus (P) deficiency. Affected palms are stunted and the stem develops with a pyramid shape.....	360
8	(a) Mid-crown yellowing (potassium [K] deficiency)	361
	(b) Confluent orange spotting (COS) due to potassium (K) deficiency is usually found on the leaflets of older palms, since K is translocated from older to younger leaves.....	361
	(c) Magnesium deficiency is characterized by the yellowing of older fronds, since Mg is translocated to younger fronds	361
9	(a) Boron deficiency symptoms may differ from one palm to another, and often from one site to the next. In all cases, the foliage is generally damaged.....	362
	(b) In copper (Cu) deficient palms, small chlorotic streaks affect the distal section of leaflets on the youngest leaves.....	362
	(c) Chlorosis due to iron (Fe) deficiency is marked by discoloration of the very last 3–4 fronds after emergence	362
10	(a) White stripes develop parallel with the midrib. Pencil-shaped discoloration extend over the whole length of the lamina either side of the leaflet midrib.....	363
	(b) Seedlings that have been correctly identified as runts should be destroyed immediately. Do not keep runts back in 'hospitals' in the hope that they will recover	363

	<i>(c) Other abnormalities – genetic origin</i>	363
11	<i>Other abnormalities – pathological origin</i>	364
12	<i>(a) Canopy management</i>	365
	<i>(b) Fertilizer application</i>	365
13	<i>Nursery selection:</i>	
	<i>A Healthy prenursery seedling</i>	366
	<i>B Healthy main nursery seedling</i>	
	<i>C Twisted shoot caused by planting the seed upside down</i>	366
	<i>D Collante (puckered fronds)</i>	366
	<i>E Narrow (grass) leaf</i>	366
	<i>F Crinkled leaf</i>	366
	<i>G Rolled leaf</i>	366
	<i>H Dwarf seedling (right) compared with a normal seedling (left)</i>	366
	<i>I Upright seedling</i>	366
14	<i>Nursery selection:</i>	
	<i>A Flat top seedling</i>	366
	<i>B Juvenile seedling</i>	366
	<i>C Short internode seedling</i>	366
	<i>D Wide internode seedling</i>	366
	<i>E Narrow pinnae seedling</i>	366
	<i>F Chimera seedling</i>	366
	<i>G Normal (top) and giant (raksaksa) seedling (bottom)</i>	366
	<i>H Crown disease</i>	366
	<i>I Herbicide damage</i>	366
15	<i>Clonal palms show much greater uniformity than D x P material</i>	368